HEDIS[®] & QARR Provider Reference Guide - Measurement Year 2025





Table of Contents

| Glossary of Terms | |
|---|----|
| Effectiveness of Care – Prevention and Screening Measures | 7 |
| Weight Assessment & Counseling for Nutrition and Physical Activity for Children and Adolescents (WCC) | 7 |
| Childhood Immunization Status (CIS) | 7 |
| Immunization Status for Adolescents (IMA) | 7 |
| Lead Screening in Children (LSC) | 7 |
| Breast Cancer Screening (BCS-E) | 8 |
| Documented Assessment after Mammogram (DBM-E) | 8 |
| Follow-Up After Abnormal Mammogram Assessment (FMA-E) | 8 |
| Cervical Cancer Screening (CCS) | 9 |
| Colorectal Cancer Screening (COL-E) | 9 |
| Chlamydia Screening in Women (CHL) | 10 |
| Care of Older Adults (COA) | 10 |
| Oral Evaluation, Dental Services (OED) | 10 |
| Topical Fluoride for Children (TFC) | 10 |
| Medical Assistance with Smoking and Tobacco Cessation (MSC) | 10 |
| COVID-19 Immunization Status (CVS) | 10 |
| Effectiveness of Care – Respiratory Conditions Measures | 11 |
| Appropriate Testing for Pharyngitis (CWP) | 11 |
| Pharmacotherapy Management of COPD Exacerbation (PCE) | 11 |
| Asthma Medication Ratio (AMR) | 12 |
| Effectiveness of Care – Cardiovascular Conditions Measures | 13 |

| Healthcare Effectiveness Data and Information Set (HEDIS®) & Quality Assurance Reporting Requirements (QARR) Provider Reference Guide - MY2025 Controlling High Blood Pressure (CBP) | 13 |
|---|----|
| Blood Pressure Control for Patients with Hypertension (BPC-E) | |
| Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) | 14 |
| Statin Therapy for Patients with Cardiovascular Disease (SPC) | 15 |
| Cardiac Rehabilitation (CRE) | 16 |
| Effectiveness of Care – Diabetes Measures | 17 |
| Glycemic Status Assessment for Patients with Diabetes (GSD) | 17 |
| Blood Pressure Control for Patients with Diabetes (BPD) | 17 |
| Eye Exam for Patients with Diabetes (EED) | 17 |
| Kidney Health Evaluation for Patients with Diabetes (KED) | |
| Statin Therapy for Patients with Diabetes (SPD) | 19 |
| Effectiveness of Care – Musculoskeletal Conditions Measures | 20 |
| Osteoporosis Management in Women Who Had a Fracture (OMW) | 20 |
| Osteoporosis Screening in Older Women (OSW) | 20 |
| Effectiveness of Care – Behavioral Health Measures | 21 |
| Diagnoses of Mental Health Disorders (DMH) | 21 |
| Follow-Up Care for Children Prescribed ADHD Medications (ADD-E) | 21 |
| Follow-Up After Hospitalization for Mental Illness (FUH) | 21 |
| Follow-Up After Emergency Department Visit for Mental Illness (FUM) | 22 |
| Diagnosed Substance Use Disorders (DSU) | 22 |
| Follow-Up After High-Intensity Care for Substance Use Disorder (FUI) | 22 |
| Follow-Up After Emergency Department Visit for Substance Use (FUA) | 23 |
| Pharmacotherapy for Opioid Use Disorder (POD) | 23 |
| Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD) | 23 |
| Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD) | 23 |

| Healthcare Effectiveness Data and Information Set (HEDIS [®]) & Quality Assurance Reporting Requirements (QARR) Provider Reference Guide - MY2025 Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC) | 24 |
|---|----|
| Adherence to Antipsychotic Medications for Individuals with Schizophrenia (SAA) | 24 |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM-E) | 24 |
| Depression Screening and Follow-Up for Adolescents and Adults (DSF-E) | 25 |
| Utilization of the PHQ-9 to Monitor Depression Symptoms for Adolescents and Adults (DMS-E) | 25 |
| Depression Remission or Response for Adolescents and Adults (DRR-E) | 25 |
| Effectiveness of Care – Care Coordination Measures | 26 |
| Advanced Care Planning (ACP) | 26 |
| Transitions of Care (TRC) | 26 |
| Follow-Up After Emergency Department Visit for People with Multiple High-Risk Chronic Conditions (FMC) | 26 |
| HIV Care | 27 |
| Viral Load Suppression (VLS) | 27 |
| Sexually Transmitted Diseases: Chlamydia Screening | 27 |
| Sexually Transmitted Diseases: Gonorrhea Screening | 27 |
| Sexually Transmitted Diseases: Syphilis Screening | 27 |
| Health Plan Descriptive Information | 28 |
| Social Needs Screening and Intervention (SNS-E) | |
| Language Diversity of Membership (LDM) | |
| Race/Ethnicity Diversity of Membership (RDM) | 28 |
| Access/Availability of Care Measures | 29 |
| Adults' Access to Preventative/Ambulatory Health Services (AAP) | 29 |
| Initiation and Engagement of Substance Use Disorder Treatment (IET) | 29 |
| Prenatal and Postpartum Care (PPC) | 29 |
| Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP) | 29 |
| Measures Collected Through Health Outcomes Survey | 30 |

| Healthcare Effectiveness Data and Information Set (HEDIS®) & Quality Assurance Reporting Requirements (QARR) Provider Reference Guide - MY2025 | |
|--|----|
| Medicare Health Outcomes Survey (HOS) | |
| Fall Risk Management (FRM) | 30 |
| Management of Urinary Incontinence in Older Adults (MUI) | |
| Physical Activity in Older Adults (PAO) | 30 |
| Utilization and Risk Adjusted Utilization Measures | 31 |
| Well-Child Visits in the First 30 Months of Life (W30) | |
| Child and Adolescent Well-Care Visits (WCV) | 31 |
| Plan All-Cause Readmissions (PCR) | 31 |
| Hospitalization Following Discharge from A Skilled Nursing Facility (HFS) | |
| Acute Hospitalization Utilization (AHU) | |
| Emergency Department Utilization (EDU) | 31 |
| Hospitalization for Potentially Preventable Complications (HPC) | 32 |
| Emergency Department Visits for Hypoglycemia in Older Adults with Diabetes (EDH) | 32 |
| References and Comments | 33 |

Glossary of Terms:

Measure Name: The name of the HEDIS® measure used for reporting

Measure Code: The code used to identify the measure name

Age Criteria: The age of the population that is eligible for the measure

Measure Description: The description of what is being measured in each domain

ICD10 Codes: The diagnosis code assigned to conditions and diseases used for claims reporting

CPT Codes: Codes used to describe the treatment being provided

Medication List: List of medications from the current year HEDIS® specifications that meet numerator compliance for the attributed measure

HCPCS Codes: Codes used to describe supplies, equipment and devices being provided

LOINC Codes: Codes use to describe the tests being provided, such as laboratory tests and vital signs

Exclusions: Some measures remove eligible members from a measure they are otherwise eligible for (ex: hospice, palliative care, ESRD/dialysis)

Measurement Year (MY): Refers to the current calendar year (January 1, 2025 – December 31, 2025), for the measures unless otherwise specified

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--|--------------------|---|---|------------------------------|
| Weight Assessment & Counseling for Nutrition and Physical Activity for Children & Adolescents (WCC) | of age | Eligible members with an outpatient PCP and/or OB/GYN visit with documentation of BMI profile and counseling for nutrition and physical activity, within the MY. | ICD 10 Codes: Z68.51-Z68.54, Z71.82, Z02.5 CPT Codes: 97802-97804 Exclusion: Members on/using hospice services (99377) members with a pregnancy diagnosis during the MY, and members who expired during the MY. | SelectHealth |
| Childhood Immunizations Status (CIS-E) | 2 years of age | Children who have received four doses of DTaP (diphtheria, tetanus, and pertussis) and PCV (pneumococcal conjugate); three doses of IPV (polio), HepB (hepatitis B), and HIB (haemophilus influenza B), two to three doses of RV (rotavirus), two Flu vaccines (influenza), and one dose of MMR (measles, mumps, and rubella), VZV (chicken pox) and HepA (Hepatitis A) by their second birthday. | ICD 10 Codes: NA CPT Codes: 90697, 90698, 90670, 90644, 90680/90681, 90655, 90674, 90707, 90710, 90633 Exclusion: Members on/using hospice services (99377), members who expired during the MY, members where a contraindication to a childhood vaccine has been identified on or before their second birthday and members who had organ and bone marrow transplants. | SelectHealth |
| Immunizations for Adolescents (IMA-E) | 13 years of age | Adolescents who have received one dose of Tdap (tetanus, diphtheria, and pertussis) and meningococcal vaccines, and have had the HPV (human papillomavirus) vaccine series finalized, by their 13 th birthday. | ICD 10 Codes: NA CPT Codes: 90649-90651, 90619, 90715 Exclusions: Members on/using hospice services (99377) and members who expired during the MY. | SelectHealth |
| Lead Screening in Children (LSC) | 2 years of age | Children who have completed at least one lead blood test (capillary or venous) by their second birthday, to determine lead poisoning. | ICD 10 Codes: NA CPT Codes: 83655 LOINC Codes: 10368-9, 10912-4, 17052-2, 5674-7 Exclusions: Members on/using hospice Services (99377) and members who expired during the MY. | SelectHealth |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--|--------------------------|--|---|--|
| Breast Cancer Screening (BCS-E) | 50 to 74 years of age | Members between 50 to 74 years of age, who have had a mammogram October 1 st two years prior through the end of the MY, to screen for breast cancer. *NCQA implemented an electronic version of this measure. Providers are encouraged to submit data administratively. | ICD 10 Codes: NA CPT Codes: 77061-77063, 77065-77067 Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who have expired during the MY, and those who have either a unilateral (Z90.11 or Z90.12) or bilateral mastectomy (Z90.13), members 66 and older residing in a facility or members 66 and older with a diagnosis of frailty (L89.000) <i>and</i> advanced illness (A81.00), and members who had gender-affirming chest surgery with a gender dysphoria diagnosis (F64.1, F64.8, F64.9). | |
| Documented Assessment After Mammogram (DBM-E) | 40 to 74 years of age | Members between 40 to74 years of age, with a BI- RADS assessment documented within 14 days of a completed mammogram. *NCQA implemented an electronic version of this measure. Providers are encouraged to submit data administratively | ICD 10 Codes: NA CPT Codes: 77061-77063, 77065-77067 RadLex Radiology Lexicon: RID36028-RID36036 Exclusions: Members on/using hospice services (99377) or members who have expired during the MY | SelectHealth, Total, EasyCare and EasyCare Plus |
| Follow-Up After Abnormal Mammogram Assessment (FMA-E) | 40 to 74 years of age | Members between 40 to 74 years of age, with inconclusive or high-risk BI-RADS assessments who received follow-up care within 90 days of a completed mammogram. *NCQA implemented an electronic version of this measure. Providers are encouraged to submit data | ICD 10 Codes: NA CPT Codes: 19081, 19083, 19085, 19100, 19101,76641- 76642, 77061-77063, 77065-77067 Exclusions: Members on/using hospice services or members who have expired during the MY | SelectHealth, Total, EasyCare and EasyCare Plus |

administratively

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|---|--------------------------|--|---|------------------------------|
| Cervical Cancer Screening (CCS-E) | 21 to 64 years of age | Members 21 to 64 years of age, who have had a cervical cytology test (21 to 64) within 3 years, cervical high-risk human papillomavirus test (30- 64) within 5 years or both cytology/hrHPV co- testing (30-64) within the last 5 years, to screen for cervical cancer. *NCQA implemented an electronic version of this measure. Providers are encouraged to submit data administratively. | ICD 10 Codes: NA CPT Codes: 88141-88143, 87624, 87625 Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, or have had a hysterectomy with no residual cervix (57530) and/or absence of cervix (Q51.5) and members assigned as male sex at time of birth. | SelectHealth |
| Colorectal Cancer Screening (COL-E) | 45 to 75 years of age | Members 45 to 75 years of age, who have had at least one of the following cancer screenings annually or within specified timeframe from the MY: Fecal Occult Blood Test (FOBT) or Fecal Immunochemical Test (FIT) annually, Stool DNA with FIT testing within 3 years, Flexible Sigmoidoscopy or CT Colonography within 5 years and/or Colonoscopy within 10 years. *NCQA implemented an electronic version of this measure. Providers are encouraged to submit data administratively. | ICD 10 Codes: NA CPT Codes: 82270, 81528, 45330, 74261, 44388 Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, members 66 and older residing in a facility or members 66 and older with a diagnosis of frailty (L89.000) <i>and</i> advanced illness (A81.00), and members with a diagnosis of colorectal cancer (C18.9) or total colectomy (44150). | |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--|-------------------------------------|---|--|--|
| Chlamydia Screening (CHL) | 16 to 24 years of age | Sexually active members between 16 to 24 years of age, who have at least one chlamydia screening test within the MY. | ICD 10 Codes: NA CPT Codes: 87110, 87320, 87490-87492 Exclusions: Members on/using hospice services (99377), members who have expired during the MY and members assigned as male sex at time of birth. | SelectHealth |
| Care of Older Adults (COA) | 66 years of age | Members who are 66 years of age who have completed both functional status assessments, as well as medication review (including OTC and supplements) at least once within the MY. | ICD 10 Codes: NA CPT Codes: 99483, 90863, 99495,1170F, 1159F & 1160F HCPCS Codes: G8427, G0438 Exclusions: Members on/using hospice services (99377), members who expired during the MY, and services provided in an acute inpatient setting (99221). | Total, EasyCare and EasyCare Plus |
| Oral Evaluation, Dental Services (OED) | Members under 21 years of age | Members under 21 years of age, who have completed a periodic or comprehensive oral exam with a dental provider during the MY. | ICD 10 Codes: NA CPT Codes: NA Exclusions: Members on/using hospice services (99377), or members who have expired during the MY. | SelectHealth |
| Topical Fluoride for Children (TFC) | 1 to 4 years of age | Members between 1 and 4 years of age, who have had two or more fluoride varnish applications completed during the MY. | ICD 10 Codes: NA CPT Codes: 99188 Exclusions: Members on/using hospice services (99377), or members who have expired during the MY. | SelectHealth |
| Medical Assistance with Smoking and Tobacco Cessation (MSC) | 18 years of age and older | This is a patient survey (CAHPS) based measure, focusing on advising smokers and tobacco users to quit, as well as discussing cessation strategies and medications. | ICD 10 Codes: NA CPT Codes: NA Exclusions: NA | SelectHealth, Total, EasyCare and EasyCare Plus |
| COVID-19 Immunization Status (CVS) | 6 months of age and older | Members 6 months of age and older, who have received at least the first dose of the COVID-19 vaccine. | ICD 10 Codes: NA CPT Codes: 91318-91322, 91304 Exclusions: Dually enrolled members in Medicaid and Medicare during the MY. | SelectHealth |

Effectiveness of Care- Respiratory Conditions Measures:

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--|--------------------------------|--|---|--|
| Appropriate Testing for Pharyngitis (CWP) | 3 years of age and older | Members 3 years of age and older, who were evaluated and diagnosed with pharyngitis, given an antibiotic and received a strep test (group A streptococcus), for each encounter. | ICD 10 Codes: J02.0, J02.8, J03.01 CPT Codes: 87070, 87081, 87430 Exclusions: Members on/using hospice services (99377), and members who expired during the MY. | SelectHealth, Total, EasyCare and EasyCare Plus |
| Pharmacotherapy Management of COPD Exacerbation (PCE) | 40 years of age | Members 40 years of age, which were hospitalized or had an emergency room visit between January 1 st and November 30 th , with COPD exacerbation, who were prescribed appropriate medications (a systemic corticosteroid or a bronchodilator). | ICD 10 Codes: J44.0, J44.9, J43.0, J43.9 Medication List: Cortisone, Dexamethasone, Hydrocortisone, Methylprednisolone, Prednisolone, Prednisone, Aclidinium bromide, Ipratropium, Tiotropium, Umeclidinium, Albuterol, Aformoterol, Formoterol, Indacaterol, Levalbuterol, Metaproterenol, Olodaterol, Salmeterol, Albuterol-ipratropium, Budesonide-formoterol, Fluticasone-salmeterol, Fluticasone-vilanterol, Fluticasone furoate- umeclidinium-vilanterol, Formoterol-aclidinium, Formoterol-glycopyrrolate, Formoterol-mometasone, Glycopyrrolate-indacaterol, Olodaterol-tiotropium and Umeclidinium-vilanterol. Exclusions: Members on/using hospice services (99377), members who have expired during the MY, and nonacute hospitalizations. | SelectHealth, Total, EasyCare and EasyCare Plus |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|-------------------------------------|-------------------------|--|---|------------------------------|
| Asthma Medication Ratio (AMR) | 5 to 64 years of age | Members 5 to 64 years of age, with a diagnosis of persistent asthma that have a controller medications vs total asthma medications prescribed ratio of at least 0.50. | ICD 10 Codes: J45.21, J45.51, J45.991 Medication List: Omalizumab, Dupilumab, Benralizumab, Mepolizumab, Reslizumab, Budesonide- formoterol, Fluticasone-salmeterol, Fluticasone- vilanterol, Formoterol- mometasone, Beclomethasone, Budesonide, Ciclesonide, Flunisolide, Fluticasone, Mometasone, Montelukast, Zafirlukast, Zileuton and Theophylline. Exclusions: Members on/using hospice services (99377), members who have expired during the MY, members who had a diagnosis of emphysema (J43.9), COPD (J44.0), chronic respiratory conditions (J68.4), cystic fibrosis (E84.9) and acute respiratory failure (J96.00), and members without any asthma controlling medications dispensed during the MY. | SelectHealth |

Effectiveness of Care- Cardiovascular Conditions Measures:

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--|--------------------------|--|---|--|
| Controlling High Blood Pressure (CBP) | 18 to 85 years of age | Members 18 to 85 years of age with a diagnosis of HTN (hypertension) whose most recent blood pressure measurement during the MY, shows an adequate control of condition, as evidenced by results of<140/90mm Hg. | ICD 10 Codes: I10 CPTII Codes: 3079F, 3078F, 3080F, 3074F, 3075F, 3077F Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who have expired during the MY, members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) and advanced illness (A81.00) diagnosis, and members 81 and older with at least two indications of frailty (L89.000) during the MY, members with a diagnosis of End-stage renal disease (ESRD) (N18.6), nephrectomy (50220 total or 50240 partial) or kidney transplant (50360), and members who are pregnant (O00.0). | SelectHealth, Total, EasyCare and EasyCare Plus |
| Blood pressure control for Patients with Hypertension (BPC-E) | 18 to 85 years of age | Members 18 to 85 years of age with a diagnosis of HTN (hypertension) whose most recent blood pressure measurement during the MY, shows an adequate control of condition, as evidenced by results of <140/90mm Hg. *NCQA implemented an electronic version of this measure. Providers are encouraged to submit data administratively | CD 10 Codes: 110 CPTII Codes: 3079F, 3078F, 3080F, 3074F, 3075F, 3077F Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who have expired during the MY, members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) <i>and</i> advanced illness (A81.00) diagnosis, and members 81 and older with at least two indications of frailty (L89.000) during the MY, members with a diagnosis of End-stage renal disease (ESRD) (N18.6), nephrectomy (50220 total or 50240 partial) or kidney transplant (50360), and members who are pregnant (O00.0). | SelectHealth, Total, EasyCare and EasyCare Plus |

| (Measure Code) ' | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|----------------------|--------------|---|---|------------------------------|
| | 18 years of | Members 18 years of age and older who were | ICD 10 Codes: I21.01, I21.19, I21.29 | SelectHealth, |
| Blocker Treatment | age and | discharged from a hospitalization for AMI (acute | Medication List: Carvedilol, Labetalol, Nadolol, Pindolol, | Total, |
| After a Heart Attack | older | myocardial infarction), between July 1 the prior | Propranolol, Timolol, Sotalol, Acebutolol, Atenolol, | EasyCare and |
| (РВН) | | year through June 30 th of the MY, that were prescribed a six-month treatment of persistent beta-blockers. | Betaxolol, Bisoprolol, Metorpolol, Nebivolol, Atenolol- chlorthalidone, Bendroflumethiazide-nadolol, Bisoprolol- hydrochlorothiazide, Hydrochlorothiazide- metroprolol and Hydrochlorothiazide-propranolol. Exclusions: Members on/using hospice services (99377), members who expired during the MY, members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) and advanced illness (A81.00) diagnosis, and members 81 and older with two or more frailty (L89.000) diagnosis, members diagnosed with asthma (J45.21), COPD (J44.0), obstructive chronic bronchitis, chronic respiratory conditions (J68.4) and hypotension (I95.9), members where it is contraindicated for beta-blocker | |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--|---|--|--|--|
| Statin Therapy for Patients with Cardiovascular Disease (SPC) | Males 21 to 75 years of age; Females 40 to 75 years of age | Male members between 21 to 75 years of age, and female members between 40 to 75 years of age, who have had an ASCVD diagnosis (atherosclerotic cardiovascular disease) and has received moderate or high- intensity statin therapy at least once during the MY, as well as maintained an 80% or higher of statin adherence. | ICD 10 Codes: I21.01, I21.9, I21.A9 High-intensity Statin Medication List: Atorvastatin 40- 80mg, Amlodipine-atorvastatin 40-80mg, Rosuvastatin 20-40mg, Simvastatin 80mg and Ezetimibe-simvastatin 80mg. Moderate-intensity Statin Medication List: Atorvastatin 10-20mg, Amlodipine-atorvastatin 10- 20mg, Rosuvastatin 5-10mg, Simvastatin 20-40mg, Ezetimibe-simvastatin 20-40mg, Pravastatin 40-80mg, Lovastatin 40mg, Fluvastatin 40-80mg and Pitavastatin 1- 4mg. Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, members 66 years of age or older residing in a facility or have a diagnosis of frailty (L89.000) <i>and</i> advanced illness (A81.00), members diagnosed with ESRD (N18.5), cirrhosis (K70.30), myalgia (M79.10), myositis (M60.80), myopathy (G72.9) or rhabdomyolysis (M62.82) caused by statins, and members with a pregnancy diagnosis (O00.0), or received IVF (in vitro fertilization) (HCPCS S4015) or prescribed clomiphene | SelectHealth, Total, EasyCare and EasyCare Plus |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--------------------------------|---------------------|--|---|-------------------------------|
| Cardiac Rehabilitation | 18 years of age and | Members 18 years of age and older, who participated in cardiac rehabilitation | ICD 10 Codes: I21.01, I21.9, I21.A9 CPT Codes: 93797, 93798 | SelectHealth, Total, |
| (CRE) | older | following an eligible cardiac event (such as MI, CABG or heart transplant). Rates reported include initiation (2 or more sessions within 30 days), engagement 1 (12 | Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) and advanced illness (A81.00) diagnosis, members 81 and older with two or more | EasyCare and EasyCare Plus |
| | | or more sessions within 90 days), engagement 2 (24 or more sessions within 180 days) and achievement (36 or more sessions within 180 days). | indications of frailty (L89.000). Members diagnosed with an MI (I21.01), CABG (33510), heart or heart & lung transplant (33927) or heart valve repair/replacement (33361) within 180 days after initial event are also excluded. | |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|---|--------------------------|---|---|--|
| Glycemic Status Assessment for Patients with Diabetes (GSD) | 18 to 75 years of age | Members between 18 to 75 years of age, diagnosed with diabetes (type 1 and 2), whose most recent glycemic status (HbA1c or GMI) was controlled (demonstrated by glycemic status <8.0) or uncontrolled (demonstrated by glycemic status >9.0) *Formerly reported as Hemoglobin A1c Control for Patients with Diabetes (HBD) | ICD 10 Codes: E10.10, E10.36, E11.00 CPT Codes: 83036, 83067 CPTII Codes: 3044F, 3051F, 3052F, 3046F Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who have expired during the MY, members 66 years of age or older residing in a facility, members 66 or older with a frailty (L89.000) <i>and</i> advanced illness (A81.00) diagnosis. | SelectHealth, Total, EasyCare and EasyCare Plus |
| Blood Pressure Control for Patients with Diabetes (BPD) | 18 to 75 years of age | Members between 18 to 75 years of age, diagnosed with diabetes (types 1 and 2), whose most recent blood pressure measurement during the MY, shows an adequate control of condition, as evidenced by results of <140/90mm Hg. | ICD 10 Codes: E10.10, E10.36, E11.00 CPTII Codes: 3079F, 3078F, 3080F, 3074F, 3075F, 3077F Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, members 66 years of age or older residing in a facility, members 66 and older with a frailty (L89.000) <i>and</i> advanced illness (A81.00) diagnosis. | SelectHealth, Total, EasyCare and EasyCare Plus |
| Eye Exam for Patients with Diabetes (EED) | 18 to 75 years of age | Members between 18 to 75 years of age, diagnosed with diabetes (types 1 and 2), who have completed a retinal eye exam. | ICD 10 Codes: E10.10, E10.36, E11.00 CPT Codes: 67028, 67030, 92201, 92230 CPTII Codes: 2022F (evidence of retinopathy), 2023F (no evidence of retinopathy), 3072F (no evidence of retinopathy in the prior year) Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, bilateral eye enucleation, members 66 years of age or older residing in a facility, members 66 and older with a frailty (L89.000) and advanced illness (A81.00) diagnosis. | SelectHealth, Total, EasyCare and EasyCare Plus |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|------------------------------------|--------------------------|--|--|-------------------------------|
| Kidney Health Evaluation for | 18 to 85 years of age | Members between 18 to 85 years of age, diagnosed with diabetes (types 1 and 2), who completed the | ICD 10 Codes: E10.10, E10.36, E11.00 CPT Codes: 82043, 82570, 80047 | SelectHealth, Total. |
| Patients with Diabetes (KED) | years of age | eGFR (estimated glomerular filtration rate) and uACR (urine albumin-creatinine ratio) lab tests during the MY. *The uACR must have both a quantitative urine albumin test and a urine creatinine test to be complete. | LOINC Codes: 13705-9, 1754-1, 2161-8, 50044-7 Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) and advanced illness (A81.00) diagnosis, members 81 and older with two indications of frailty (L89.000) diagnosis, members diagnosed with ESRD (N18.5) or on dialysis. | EasyCare and EasyCare Plus |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--|--------------------------|--|--|------------------------------|
| Statin Therapy for Patients with Diabetes (SPD) | 40 to 75 years of age | Members between 40 to 75 years of age, with a diagnosis of diabetes, who do not have an ASCVD diagnosis (atherosclerotic cardiovascular disease), and has received statin therapy, as well as maintained an 80% of statin adherence. | ICD 10 Codes: E10.10, E10.36, E11.00 High-intensity Statin Medication List: Atorvastatin 40-80mg, Amlodipine-atorvastatin 40-80mg, Rosuvastatin 20-40mg, Simvastatin 80mg and Ezetimibesimvastatin 80mg. Moderate-intensity Statin Medication List: Atorvastatin 10-20mg, Amlodipine-atorvastatin 10-20mg, Rosuvastatin 5-10mg, Simvastatin 20-40mg, Ezetimibe-simvastatin 20-40mg, Pravastatin 20-40mg, Lovastatin 40mg, Fluvastatin 40-80mg and Pitavastatin 1-4mg. Low-intensity Statin Medication List: Ezetimibesimvastatin 10-20mg and Simvastatin 5-10mg. Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, members 66 years of age or older residing in a facility, members 66 and older with a frailty (L89.000) and advanced illness (A81.00) diagnosis, members diagnosed with MI (I21.9), CABG or PCI during the MY, and members with a diagnosis of ESRD (N18.5), cirrhosis (K70.30), myalgia (M79.10), myositis (M60.80), myopathy (G72.9) or rhabdomyolysis (M62.82) caused by statin and members with a pregnancy (000.0) diagnosis, or received IVF (in vitro fertilization) HCPCS S4015) or prescribed clomiphene during the MY. | |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--|--------------------------|---|--|---|
| Osteoporosis Management in Women Who Had a Fracture (OMW) | 67 to 85 years of age | Female members between 67 to 85 years of age, who have had a BMD (bone mineral density) test or prescribed a medication for osteoporosis treatment within six months, after suffering a fracture. | ICD 10 Codes: M80.00XA, M80.80XA CPT Codes: 76977, 77078, 77080-77081 HCPCS Codes: J0897, J1740, J3110, J3111, J3489 Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, members 67 years of age or older residing in a facility, members 67 to 80 years of age with a frailty (L89.000) <i>and</i> advanced illness (A81.00) diagnosis, members 81 and older with two or more indications of frailty (L89.000) diagnosis. | Total, EasyCare and EasyCare Plus |
| Osteoporosis Screening in Older Women (OSW) | 66 to 75 years of age | Female members between 65-75 years of age, who have been assessed for osteoporosis during the MY. | ICD 10 Codes: NA CPT Codes: 76977, 77078, 77080, 77081, 77085 Exclusions: Members on/using hospice services (99377) or palliative care (Z51.5), members who expired during the MY, members 66 years of age or older with a frailty (L89.000) <i>and</i> advanced illness (A81.00) diagnosis, and members who had encounters for osteoporosis therapy anytime through December 31 st of the MY or claims for osteoporosis medications within 3 years prior to the MY. | Total, EasyCare and EasyCare Plus |

Effectiveness of Care- Musculoskeletal Conditions Measures:

Effectiveness of Care- Behavioral Health Measures:

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--|--------------------------------|---|---|--|
| Diagnoses of Mental Health Disorders (DMH) | 1 years of age and older | Members 1 years of age and older, who have been diagnosed with a mental health disorder during the MY Age stratifications are 1-17, 18-64 and 65 years of age and older. | ICD 10 Codes: F03.90, F21, F31.0, F43.10 CPT Codes: NA Exclusions: Members on/using hospice services (99377) and members who expired during the MY. | SelectHealth, Total, EasyCare and EasyCare Plus |
| Follow-up Care for Children Prescribed ADHD Medication (ADD-E) | 6 to 12 years of age | Members between 6 to 12 years of age, who were prescribed ADHD (attention deficit/hyperactivity disorder) medications and had a follow-up visit within 30 days of prescription <i>(initiation phase)</i> and at least two more follow-up visits within 10 months of prescription <i>(continuation and maintenance phase).</i> | | SelectHealth |
| Follow-Up After Hospitalization for Mental Illness (FUH) | 6 years of age and older | Members 6 years of age and older, who had a primary diagnosis of mental illness or self-harm diagnosed hospital admission, which was discharged and had a mental health provider follow-up. Follow-up appointments are rated by those who complete an appointment within 7 days post- discharge, and those who complete an appointment within 30 days post-discharge. | ICD 10 Codes: T14.91XA, F20.0, F31.0, F32.9 CPT Codes: 98960-98962, 99202-99205 Exclusions: Members on/using hospice services (99377), members who expired during the MY, and any members discharged and readmitted within 30 days. | SelectHealth, Total, EasyCare and EasyCare Plus |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--|---------------------------------|--|--|--|
| Follow-Up After Emergency Department Visit for Mental Illness (FUM) | 6 years of age and older | Members 6 years of age and older, who had a principal diagnosis of mental illness or self-harm diagnosed emergency department (ED) visit, and had a mental health provider follow-up. Follow-up appointments are rated by those who complete an appointment within 7 days post ED- discharge, and those who complete an appointment within 30 days post ED- discharge. | ICD 10 Codes: T14.91XA, F20.0, F31.0, F32.9 CPT Codes: 98960-98962, 99202-99205 Exclusions: Members on/using hospice services (99377), members who expired during the MY. | SelectHealth, Total, EasyCare and EasyCare Plus |
| Diagnosed Substance Use Disorders (DSU) | 13 years of age and older | Members 13 years of age and older with a substance use disorder during the MY. Substance abuse disorders include alcohol, opioid, substance use and other/unspecified drugs. | ICD 10 Codes: F10.10, F11.10, F14.10, F15.10 CPT Codes: NA Exclusions: Members on/using hospice services (99377) and members who expired during the MY. | SelectHealth, Total, EasyCare and EasyCare Plus |
| Follow-Up After High-Intensity Care for Substance Use Disorder (FUI) | 13 years of age and older | Members 13 years of age and older with a substance use disorder, which had an acute inpatient hospitalization, residential treatment, or detoxification visit, resulting in follow-up visits. Follow-up appointments are rated by those who complete an appointment within 7 days post- discharge, and those who complete an appointment within 30 days post-discharge. | ICD 10 Codes: F10.10, F11.10, F14.10, F15.10 CPT Codes: 98960-98962, 99202-99205 HCPCS Codes: H0017-H0019, H0007-H0020 Exclusions: Members on/using hospice services (99377), members who expired during the MY. | SelectHealth, Total, EasyCare and EasyCare Plus |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|---|------------------------------|--|---|--|
| Follow-Up After Emergency Department Visit for Substance Use (FUA) | 13 years of age and older | Members 13 years of age and older that had an emergency department visit with a primary substance use disorder or drug overdose diagnosis and completed a follow-up visit. Follow-up appointments are rated by those who complete an appointment within 7 days post-discharge, and those who complete an appointment within 30 days post-discharge. | ICD 10 Codes: T40.0X1A, F10.10, F11.10, F14.10, F15.10 CPT Codes: 98960-98962, 99202-99205 HCPCS Codes: H0020, H0004, G2075 Exclusions: Members on/using hospice services (99377), members who expired during the MY. | SelectHealth, Total, EasyCare and EasyCare Plus |
| Pharmacotherapy for Opioid Use Disorder (POD) | 16 years of age and older | Members 16 years of age and older with an opioid use disorder, who received new pharmacotherapy that lasted for at least 180 days. | ICD 10 Codes: F11.10, F11.20 CPT Codes: NA HCPCS Codes: G2070, G2069, J0572, H0033, H0020, J2315 Exclusions: Members on/using hospice services (99377), members who expired during the MY. | SelectHealth, Total, EasyCare and EasyCare Plus |
| Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD) | 18 to 64 years of age | Members between 18 and 64 years of age, diagnosed with schizophrenia, schizoaffective or bipolar disorder with a completed diabetes screening, which were prescribed antipsychotic medications during the MY. | ICD 10 Codes: F31.9, F20.9, F25.9 CPT Codes: 80047, 80053, 83036, 83037 Exclusions: Members on/using hospice services (99377), members who expired during the MY, Members who are diagnosed with diabetes, and members that did not receive antipsychotic medications during the MY. | SelectHealth |
| Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD) | 18 to 64 years of age | Members between 18 and 64 years of age, with a diagnosis of diabetes and schizophrenia or schizoaffective disorder, who completed an HbA1c and LDL-C test during the MY. | ICD 10 Codes: E10.9, E11.9, F20.9, F25.9 CPT Codes: 83036, 83037, 80061, 83700 CPT II Codes: 3044F, 3051F, 3052F, 3046F, 3048F, 3049F, 3050F Exclusions: Members on/using hospice services (99377), members who expired during the MY. | SelectHealth |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|---|------------------------------|--|---|--|
| Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC) | 18 to 64 years of age | Members between 18 and 64 years of age, with a diagnosis of cardiovascular disease and schizophrenia or schizoaffective disorder, which completed LDL-C testing during the MY. | ICD 10 Codes: I20.0, I24.9, I21.4, F20.9, F25.9 CPT Codes: 80061, 83700, 83721 CPTII Codes: 3048F, 3049F, 3050F Exclusions: Members on/using hospice services (99377) and members who expired during the MY. | SelectHealth |
| Adherence to Antipsychotic Medications for Individuals with Schizophrenia (SAA) | 18 years of age and older | Members 18 years of age and older diagnosed with schizophrenia or schizoaffective disorder, which were prescribed an antipsychotic medication, and remained on treatment for at least 80% of the period. | ICD 10 Codes: F20.3, F20.89, F20.9, F25.8, F25.9 CPT Codes: NA HCPCS Codes: J0401, J1631, J1944, J2426, J2680 Exclusions: Members on/using hospice services (99377), members who expired during the MY, members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) <i>and</i> advanced illness (A81.00) diagnosis, members 81 and older with at least two frailty (L89.000) diagnosis, members with a diagnosis of Dementia (F03.90) or do not have at least two antipsychotic prescription refill events. | SelectHealth, Total, EasyCare and EasyCare Plus |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM-E) | 1 to 17 years of age | Members between 1 and 17 years of age, who were prescribed two or more antipsychotic medications and had metabolic testing. <i>Metabolic testing includes blood glucose</i> <i>testing, cholesterol testing or both.</i> <i>*NCQA implemented an electronic version of</i> <i>this measure. Providers are encouraged to</i> <i>submit data administratively</i> | ICD 10 Codes: NA CPT Codes: 82947, 83036, 80061 CPTII Codes: 3044F, 3046F, 3051F, 3052F, 3048F, 3049F, 3050F Exclusions: Members on/using hospice services (99377) and members who expired during the MY. | SelectHealth |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|---|------------------------------|---|---|--|
| Depression Screening and Follow-Up for Adolescents and Adults (DSF-E) | 12 years of age and older | Members 12 years of age and older who were evaluated for depression utilizing standardized screening tools, and, if a positive screen, received a follow-up care within 30 days. *NCQA implemented an electronic version of this measure. Providers are encouraged to submit data administratively. | ICD 10 Codes: NA CPT Codes: 99202-99205 LOINC Codes: 44261-6, 55758-7, 89208-3, 89205-9, 90853-3, 48545-8, 71956-8 Exclusions: Members on/using hospice services (99377), members who expired during the MY, and members diagnosed with bipolar disorder (F31.30) or depression (F32.4) | SelectHealth, Total, EasyCare and EasyCare Plus |
| Utilization of the PHQ-9 to Monitor Depression Symptoms for Adolescents and Adults (DMS-E) | 12 years of age and older | submit data administratively Members 12 years of age and older, diagnosed with major depression or dysthymia, having a documented PHQ-9 screening with score present in the medical record during an outpatient visit. *NCQA implemented an electronic version of this measure. Providers are encouraged to submit data administratively | (F32.A). ICD 10 Codes: NA CPT Codes: 99202-99205 Exclusions: Members on/using hospice services (99377), members who expired during the MY, and members diagnosed with bipolar disorder (F31.30), personality disorder (F60.3), psychotic disorder (F28) or pervasive developmental disorder (F84.9). | SelectHealth, Total, EasyCare and EasyCare Plus |
| Depression Remission or Response for Adolescents and Adults (DRR-E) | 12 years of age and older | Members 12 years of age and older, diagnosed with major depression and an elevated PHQ-9 score, having supported documentation of successful remission (score <5) or response (score decreased by 50%) within 4 to 8 months following the initial high score. *NCQA implemented an electronic version of this measure. Providers are encouraged to submit data administratively | ICD 10 Codes: NA CPT Codes: 99202-99205 Exclusions: Members on/using hospice services (99377), members who expired during the MY, and members diagnosed with bipolar disorder (F31.30), personality disorder (F60.3), psychotic disorder (F28) or pervasive developmental disorder (F84.9). | SelectHealth, Total, EasyCare and EasyCare Plus |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--------------------------------|---------------|--|--|---------------------------|
| | | | | |
| Advance Care Planning | | Members between 66 and 80 years of age who | ICD 10 Codes: M62.81, Z74.01, G30.0, G30.9 | Total, EasyCare |
| (ACP) | age and older | have an advanced illness or frailty diagnosis, or | CPT Codes: 99483, 99497 | and |
| | | receiving palliative care, and members 81 years | Exclusions: Members on/using hospice services (99377) | EasyCare Plus |
| | | of age and older who have received | and members who expired during the MY. | |
| | 10 | advanced care planning in the MY. | | T + 1 F = 0 |
| Transitions of Care | 18 years of | Members 18 years of age and older, who have | ICD 10 Codes: NA | Total, EasyCare |
| (TRC) | age and older | been discharged after an in-patient | CPT Codes: 99483, 99495, 98966, 99202-99205 | and |
| | | hospitalization, and have the following noted in | Exclusions: Members on/using hospice services (99377) | EasyCare Plus |
| | | their outpatient medical record: | and members who expired during the MY. | |
| | | 1. Notification of Inpatient Admission within 2 | | |
| | | days of discharge. | | |
| | | 2. Receipt of Discharge information within 2 | | |
| | | days of discharge. | | |
| | | 3. Patient Engagement within 30 days of | | |
| | | discharge. (Cannot be completed the same day of discharge) | | |
| | | Medication Reconciliation within 30 days | | |
| | | of discharge. (It is recommended that | | |
| | | providers use CPTII code 1111F) | | |
| Follow-Up After | 18 years of | Members 18 years of age and older, diagnosed | ICD 10 Codes: J44.9, G30.9, N18.9, I50.9, I21.9, I48.91, | Total, EasyCare |
| Emergency | age and older | with multiple high-risk chronic conditions, who | 163.9 | and |
| Department Visit for | | have been in the emergency department (ED) | CPT Codes: 99495, 98966, 99202-99205, 90791 | EasyCare Plus |
| People with Multiple | | and completed a follow-up appointment within | Exclusions: Members on/using hospice services (99377) | Laby care rido |
| High-Risk Chronic | | 7 days of the ED visit. | and members who expired during the MY. | |
| Conditions | | | | |
| (FMC) | | High-risk conditions include, but not limited to, | | |
| | | COPD, Alzheimer's disease, chronic kidney | | |
| | | disease, heart failure, acute myocardial | | |
| | | infarction, atrial fibrillation, and stroke. | | |

Effectiveness of Care- Care Coordination Measures:

HIV Care

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--|-----------------------------|---|---|---------------------------|
| Viral Load Suppression (VLS) | 2 years of age and older | Members 2 years of age and older, who are diagnosed with HIV (Human Immunodeficiency Virus), with their last HIV Viral Load Testing of the year showing a viral load less than two hundred copies/mL. | ICD 10 Codes: NA CPT Codes: NA Exclusions: NA | SelectHealth |
| Sexually Transmitted Infections: Chlamydia Screening | 13 years of age and older | Members 13 years of age and older, who have completed a chlamydia screening within the MY. | ICD 10 Codes: NA CPT Codes: NA Exclusions: Members who are under 18 years of age and are not sexually active | SelectHealth |
| Sexually Transmitted Infections: Gonorrhea Screening | | Members 13 years of age and older, who have completed a gonorrhea screening within the MY. | ICD 10 Codes: NA CPT Codes: NA Exclusions: Members who are under 18 years of age and are not sexually active | SelectHealth |
| Sexually Transmitted Infections: Syphilis Screening | 13 years of age and older | Members 13 years of age and older, who have completed a serologic syphilis screening within the MY. | ICD 10 Codes: NA CPT Codes: NA Exclusions: Members who are under 18 years of age and are not sexually active | SelectHealth |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--|--------------|--|--|--|
| Social Needs Screening and Intervention (SNS-E) | All ages | All members who have been screened for Food, Housing and Transportation insecurity/needs, as well as have documented interventions within 30 days of a positive screening. Screenings should be completed at least annually. *NCQA implemented an electronic version of this measure. Providers are encouraged to submit data administratively | ICD 10 Codes: NA CPT Codes: 96156, 96160, 96161, 97802-97804 LOINC Codes: 88122-7, 95264-8, 71802-3, 96778-6, 93030-5 Exclusions: Members on/using hospice services (99377) and members who expired during the MY, and members over 66 years of age residing in a facility. | SelectHealth, Total, EasyCare and EasyCare Plus |
| Language Diversity of Membership (LDM) Race/Ethnicity Diversity of Membership (RDM) | All ages | All members who have been screened for their preferred spoken and written languages, which have been enrolled in the Health Plan at any time during the MY. Document and report the race and ethnicity of all members, enrolled in the Health Plan at any time during the MY. | ICD 10 Codes: NA CPT Codes: NA Exclusions: NA ICD 10 Codes: NA CPT Codes: NA Exclusions: NA | SelectHealth, Total, EasyCare and EasyCare Plus SelectHealth, Total, EasyCare and EasyCare Plus |

Health Plan Descriptive Information

Access/Availability of Care Measures:

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--|---------------------------------|---|--|--|
| Adults' Access to Preventative/Ambulatory Health Services (AAP) | 20 years of age and older | Members 20 years of age and older, who have had a preventative or ambulatory care visit within the MY. | CPT Codes: 99202-99205, 99345 Exclusions: Members on/using hospice services (99377) and members who expire during the MY. | SelectHealth, Total, EasyCare and EasyCare Plus |
| Initiation and Engagement of Substance Use Disorder Treatment (IET) | 13 years of age and older | Members 13 years of age and older, who have a new substance use disorder, which initiated and engaged in treatment. Initiation of SUD treatment should be completed within 14 days and engagement of SUD treatment completed within 34 days of initiation. | ICD 10 Codes: F10.10, F11.10, F15.10 CPT Codes: 99408, 98960, 99212 HCPCS Codes: G2086, G2075, G2080 Exclusions: Members on/using hospice services (99377), members who expired during the MY. | SelectHealth, Total, EasyCare and EasyCare Plus |
| Prenatal and Postpartum Care (PPC) | None specified | Members who have had live births on October 8 th the year prior through October 7 th of the MY. Measure includes timeliness of prenatal care during the first trimester, as well as postpartum care between 7 and 84 days post-delivery. <i>Members who have multiple live births during one</i> <i>pregnancy count once, however multiple</i> <i>pregnancies resulting in two separate live</i> <i>deliveries are counted twice.</i> | ICD 10 Codes: 026.90, 026.91, 026.92, 026.93 CPT Codes: 99202-99205, 99500, 59400, 57170, 58300, 98966-98968 Exclusions: Members on/using hospice services (99377), members who expired during the MY. | SelectHealth |
| Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP) | 1 to 17 years of age | Members between 1 and 17 years of age who were prescribed a new antipsychotic medication, with documented first-line psycho-social care. | ICD 10 Codes: F20.9, F25.9, F33.3, F22 CPT Codes: 90832-90834, 90836-90840 Exclusions: Members on/using hospice services (99377), members who expired during the MY and members with antipsychotic medication as first-line care. | SelectHealth |

| Measures Collected Through the Health Outcomes Surv | ey: |
|--|-----|
|--|-----|

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--|------------------|--|--|---------------------------|
| Medicare Health Outcomes | | This survey provides an idea of how well a | ICD 10 Codes: NA | Total, EasyCare |
| Survey | age and | Medicare Advantage Organization manages | CPT Codes: NA | and |
| (HOS) | older | their members mental and physical health | Exclusions: Members on/using hospice services (99377) | EasyCare Plus |
| | | over a two year period. The rates of health- | per CMS administrative records. | |
| | | status are indicated as better, same and | | |
| | | worse than expected. | | |
| Fall Risk Management | 65 years of | Members who are 65 years of age and older, | ICD 10 Codes: NA | Total, EasyCare |
| (FRM) | age and | who have seen a practitioner in the past 12 | CPT Codes: NA | and |
| | older | months and have discussed falls or | Exclusions: Members on/using hospice services (99377) | EasyCare Plus |
| | | gait/balance problems, and those who have | per CMS administrative records. | |
| | | fallen or had gait/balance problems in the | | |
| | | last month that have received education on | | |
| | | prevention and treatment from their | | |
| Management of this are | CE | practitioner. | | Tatal Familian |
| Management of Urinary Incontinence in Older | 65 years of | Members who are 65 years of age and older, | ICD 10 Codes: NA | Total, EasyCare |
| Adults | age and older | that have reported urine leakage problems | CPT Codes: NA | and |
| | older | with their health care providers in the past 6 months, who have discussed the following | Exclusions: Members on/using hospice services (99377) per CMS administrative records. | EasyCare Plus |
| (MUI) | | with their health care provider: urinary | per civis automistrative records. | |
| | | incontinence, available treatment choices | | |
| | | and the impact urinary incontinence had on | | |
| | | daily living and sleep. | | |
| Physical Activity in Older | 65 years of | Members who are 65 years of age and older, | ICD 10 Codes: NA | Total, EasyCare |
| Adults | age and | who have seen their doctor and discussed | CPT Codes: NA | and |
| (PAO) | older | their physical activity level, and those who | Exclusions: Members on/using hospice services (99377) | EasyCare Plus |
| () | | received advice on starting, increasing or | per CMS administrative records. | |
| | | maintaining it, during the MY. | | |

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|---|---------------------------------|---|--|---|
| Well-Child Visits in the First 30 Months of Life (W30) | 15 to 30 months of age | Members between 31 days old to 15 months of age, who have had 6 or more well-child PCP visits within 15 months, as well as 2 or more well-child PCP visits between 15-30 months during the MY. | ICD 10 Codes: Z00.00, Z00.121, Z00.2, Z00.3 CPT Codes: 99381-99385, 99391-99395 Exclusions: Members on/using hospice services (99377) and members who expire during the MY. | SelectHealth |
| Child and Adolescent Well-Care Visits (WCV) | 3 to 21 years of age | Members between 3 and 21 years of age, who have had one or more well-visits with their PCP or OB/GYN during the MY. | ICD 10 Codes: Z00.00, Z00.121, Z00.2, Z00.3 CPT Codes: 99381-99385, 99391-99395 Exclusions: Members on/using hospice services (99377) and members who expire during the MY. | SelectHealth |
| Plan All-Cause Readmissions (PCR) | 18 years of age and older | Members who are 18 years of age and older, that have had an unplanned readmission, within 30 days of an observation event or inpatient admission during the MY. *Readmission can be for any diagnosis | ICD 10 Codes: NA CPT Codes: 99221-99223, 99211-99215 Exclusions: Members on/using hospice services (99377). | SelectHealth, Total, EasyCare, and EasyCare Plus |
| Hospitalization Following Discharge from a Skilled Nursing Facility (HFS) | 65 years of age and older | Members who are 65 years of age and older, who have transitioned from a Skilled Nursing Facility to the community and have had an unplanned inpatient admission within 30 to 60 days post transition. | ICD 10 Codes: NA CPT Codes: 99221-99223, 99211-99215 Exclusions: Members on/using hospice services (99377) and members residing in an institution long-term in the MY. | Total, EasyCare and EasyCare Plus |
| Acute Hospitalization Utilization (AHU) | 18 years of age and older | The ratio for risk-adjustment of observed to expected inpatient admissions and observation discharges during the MY, for members who are 18 years of age and older. | ICD 10 Codes: NA CPT Codes: 99221-99223, 99211-99215 Exclusions: Members on/using hospice services (99377). | SelectHealth, Total, EasyCare, and EasyCare Plus |
| Emergency Department Utilization (EDU) | 18 years of age and older | The ratio for risk-adjustment of observed to expected emergency department visits during the MY, for members who are 18 years of age and older. | ICD 10 Codes: NA CPT Codes: 99281-99285 Exclusions: Members on/using hospice services (99377), members with encounters for diagnosis of mental health or chemical dependency, electroconvulsive therapy and psychiatry. | Total, EasyCare and EasyCare Plus |

Utilization and Risk Adjusted Utilization Measures:

| Measure Name (Measure Code) | Age Criteria | Measure Description | ICD 10 Codes, CPT Codes and Exclusions | Line of Business (LOB) |
|--|---------------------------------|---|--|---|
| Hospitalization for Potentially Preventable Complications (HPC) | 67 years of age and older | Discharge rates for ambulatory care sensitive conditions per 1,000 members and the ratio for risk-adjustment of observed to expected discharges for acute and chronic ambulatory care sensitive conditions, for members who are 67 years of age and older. | ICD 10 Codes: NA CPT Codes: 99221-99223, 99304-9909 Exclusions: Members on/using hospice services (99377), members residing in an institution long-term, and members enrolled in an institutional SNP during the MY. | Total, EasyCare and EasyCare Plus |
| Emergency Department Visits for Hypoglycemia in Older Adults with Diabetes (EDH) | - | The ratio for risk-adjustment of observed to expected emergency department visits related to hypoglycemia during the MY. Rates are reported for members 67 years of age and older with types 1 and 2 Diabetes, with a subset of those with one or more insulin dispensing events within a 6-month treatment period July 1 the prior year through December 31 of the MY. | ICD 10 Codes: E10.641, E11.641, E13.641, E16.2 CPT Codes: 99281-99285 Exclusions: Members on/using hospice services (99377). | Total, EasyCare and EasyCare Plus |

References and Comments:

- HEDIS [®] is a registered trademark of the National Committee for Quality Assurance (NCQA). Refer to <u>www.ncqa.org/hedis/</u>
- CPT[®] (Current Procedural Terminology) is a registered trademark of the American Medical Association
- HCPCS: Healthcare Common Coding Procedure Coding System Level II
- LOINC: Logical Observation Identifiers Names and Codes
- ICD-10: Centers for Medicare & Medicaid Services at <u>www.cms.gov/Medicare/Coding/icd10/</u>
- For additional information on Advanced Illness and Frailty, refer to https://www.ncqa.org/blog/improving-care-advanced-illness-frailty/
- Quality Assurance Reporting Requirements (QARR) are adopted from NCQA HEDIS[®] with New York state-specific measures added to address health issues of particular importance in New York. This document is not inclusive of all codes and measures. The measure descriptions and codes in this document are derived from the MY2025 HEDIS[®] Technical Specifications/Value Set Directory, and the Quality Assurance Reporting Requirements Technical Specifications manual. For the full list of codes, please refer to the NCQA store to purchase.



Have questions?

Please contact us at <u>QualityManagement@vnshealth.org</u> or <u>Provider_Education@vnshealth.org</u>

For additional information and resources, please visit our provider website and toolkit at <u>https://www.vnshealthplans.org/for-health-professionals-overview/</u>